arXiv:1110.0783v2 [hep-ph] 20 Mar 2012

DESY 11-172

Violations of Lorentz invariance in the neutrino sector after OPERA

Luca Maccione,[1, *] Stefano Liberati,[2, 3, †] and David M. Mattingly[4, ‡]
[1]*DESY, Theory Group, Notkestraße 85, D-22607 Hamburg, Germany*
[2]*SISSA, Via Bonomea 265, 34136, Trieste, Italy*
[3]*INFN, Sezione di Trieste, Via Valerio, 2, 34127, Trieste, Italy*
[4]*Department of Physics, University of New Hampshire, Durham, NH 03824, USA*

The OPERA collaboration has recently reported that neutrinos travel faster than light. We review the theoretical situation of constraints on violations of Lorentz invariance, focusing in particular on the compatibility between the OPERA results with both previous constraints and ones derived after OPERA. We generalize to higher order operators the recent constraint provided by the absence of neutrino energy loss, via electron-positron pair production at OPERA energies, and show that simple modified dispersion relations in vacuo within an effective field theory framework cannot explain OPERA measurements.

PACS numbers:

I. INTRODUCTION

Lorentz symmetry is one of the grounding pillars of special relativity and hence our present understanding of the physical world. As such, its validity has been checked by numerous experiments over the last century. Despite no experimental indications of any deviations, some approaches to quantum gravity suggest that due to the fundamental structure of spacetime Lorentz symmetry might not be an exact symmetry of nature but merely an approximate or emergent symmetry at low energies that is strongly broken above some energy scale. Searches for any violation of Lorentz symmetry (LIV) have therefore received more and more attention over the last few years as possible signals of physics beyond the standard model.

Independent from any theoretical motivation, experimental constraints on LIV are very tight and so any experimental indication of a deviation from Lorentz invariance is of high importance. Recently, the OPERA collaboration reported [1] a measurement of the time of arrival of non-oscillated ν_μ (with a contamination of $\bar{\nu}_\mu$, $\bar{\nu}_e$ and ν_e that is estimated to be less than 5%) over the path length ~ 730 km from CERN to Gran Sasso with average energy of ~ 17 GeV. The ν_μ arrived earlier than light would, by an amount

$$\Delta t = 60.7 \pm 6.9\,(\mathrm{stat.}) \pm 7.4\,(\mathrm{sys.})\ . \tag{1}$$

No hints of energy or seasonal/sidereal dependence were found over the energy range of the experiment. This measurement can be converted into a measure of the propagation speed of the ν_μ's, which yields a propagation velocity for non-oscillated ν_μ [1] of

$$\frac{\Delta c}{c} = (2.48 \pm 0.28\,(\mathrm{stat.}) \pm 0.30\,(\mathrm{sys.})) \times 10^{-5}\ , \tag{2}$$

where c is the low energy speed of light in vacuum, and $\Delta c = v_\nu - c$. [1] Previously, the MINOS Collaboration reported [3] $\Delta c/c \lesssim (5.1 \pm 2.9) \times 10^{-5}$ for 3 GeV ν_μ, compatible with 0 within 2σ.

Without additional new physics, the OPERA result is a statement about the behavior of (almost) free propagating neutrinos. The relevant modification to the neutrino Lagrangian sensitive to these results is a modification to the free field kinetic term, or equivalently the particle's dispersion relation. Below we parametrize, using some simplifying

---

[*] Electronic address: luca.maccione@desy.de
[†] Electronic address: liberati@sissa.it
[‡] Electronic address: dyo7@unh.edu

[1] A few months after the publication of this paper, the Opera collaboration found two possible sources of error in their experiment. The two new effects shift the time of flight result in opposite directions and at this point there is no new data release by Opera. However, the Icarus experiment reported recently [2] on a re-analysis of the most recent data taken by Opera. Icarus finds a neutrino propagation velocity compatible with speed of light within the experimental uncertainties. Since our work does not attempt to explain the Opera claim but merely elucidates the difficulties the result has with conservative modifications of the Standard Model we shall keep the discussion basically unchanged in what follows.

assumptions and previous experimental results, a commonly discussed class of LIV effects on free field propagation within an effective field theory (EFT) framework and explain where each of the possible modifications runs into experimental constraints. The advantage of this procedure, with respect to other purely kinematical/phenomenological approaches, is that the structure of the SM is preserved, and hence observational consequences are completely calculable. Furthermore, the simplified EFT framework here adopted allows a transparent physical interpretation of the constraints casted on the salient parameters characterizing the departure from Lorentz invariance.

Since no sidereal variations have been measured for OPERA (or any other experiment) there is no experimental reason we cannot assume that the dispersion relations for neutrinos are rotationally symmetric. We do this for simplicity, although a more complicated LIV pattern could be chosen. Such an assumption picks out a preferred frame, and the natural preferred frame is that of the cosmic microwave background. Since the boost factor of an Earth centered frame is only $\sim 10^{-3}$ [4] with respect to the CMB frame, any rotation breaking effects generated by the Earth's relative motion with respect to the CMB will be naturally suppressed by a factor of $\sim 10^{-3}$ relative to rotationally invariant effects.

In short, the modified dispersion relation that we consider for a particle of species i in the following is

$$E^2 = p^2 + m_i^2 + \xi_i \frac{p^n}{M_i^{n-2}} \, , \tag{3}$$

with E and p being the energy and the momentum of the particle respectively, m the particle mass, $\xi_i = \pm$ and M_i the energy scale at which LIV appears. It is, of course, possible that the eigenstates of the LIV terms do not correspond to the energy eigenstates for neutrinos. We will not analyze the OPERA results of such a case in detail here since, as we shall see, existing neutrino constraints are so tight that all neutrino states must have essentially identical dispersion relations thereby rendering flavor dependent dispersion an unnecessary complication. The dispersion relation Eq. (3) has also been often considered in the literature as a test model for how quantum gravitational effects might influence infrared physics [5–8]. For an exhaustive classification of neutrino operators that can be added to the standard model see [9, 10].

We leave the index n as a free phenomenological parameter and consider the cases $n = 2, 3, 4$. The case $n = 1$ would produce huge effects at low energy and is strongly constrained. We shall show that due to the strong energy dependence $n = 4$ is in conflict with the OPERA results and so any $n > 4$ is also ruled out. LIV enters only through CPT odd operators at $n = 3$, while CPT invariance is preserved if $n = 2$ or $n = 4$ (while CPT breaking implies Lorentz breaking in any local EFT – see [11] – the reverse is not true and Lorenz violation does not imply CPT violation). Therefore coefficients for particles and anti-particles are related by $\xi_{\bar{i}} = (-1)^n \xi_i$ [15].

One may wonder if more general dispersion relations, in which for example various terms with different n are considered to contribute together, can be studied. However, this would require a strong fine tuning. Indeed, if the violation of special relativity comes in from new physics at some energy M, then the natural contribution of each Lorentz violating correction term changes the speed of the neutrino by a factor $c_n \, p^{n-2}/M^{n-2}$, where n is the order of the dispersion correction and c_n is a dimensionless coefficient. If M is not near OPERA energies, then the c_n's must be very carefully chosen, i.e. fine tuned, to contribute equally to the change in the neutrino speed. The alternative is to have M, the scale of the new physics, to be at OPERA energies such that the c_n can all be of the same (perhaps small) size. This would, however, be problematic as physics at 10 GeV is obviously exceptionally well explored and one would still have to account for the smallness of the c_n.

Therefore, we assume here that there is a hierarchy of terms governed by new physics at some mass scale M and check each order correction in energy, which corresponds to the assumption that M is not at 10 GeV and that the coefficients are not fine tuned, so that naturally there is a dominant term at any given neutrino energy. Without any custodial symmetry, one would expect the relevant operator that generates $n = 2$ changes in the dispersion to dominate (see e.g. [12]). We also check $n = 3$ and $n = 4$ as several theoretical models suggest this kind of dispersion relations to be the dominant terms. $n = 3$ or $n = 4$ dispersion corrections would, of course, require some other unknown physics to prevent the $n = 2$ term to be dominant (see again [12] and [13, 14] for a recent discussion of these issues).

Several constraints have been cast on LIV in different particle sectors using this approach. For example, photon dispersion is constrained by absence of birefringence to the level $M \gtrsim 10^{14} M_{\rm Pl}$ in the case $n = 3$ [16, 17], and a constraint of order $M \gtrsim 10^4 M_{\rm Pl}$ for $n = 4$ is expected from ultra-high-energy γ-rays [18]. Electron/positron dispersion relation is constrained by observations of synchrotron radiation from the Crab Nebula to a level $M \gtrsim 10^5 M_{\rm Pl}$ for $n = 3$ [19], while again ultra-high-energy cosmic ray observations help constrain the $n = 4$ case to $M \gtrsim 10^3 M_{\rm Pl}$ [18]. Also in the hadronic sector constraints coming from ultra-high-energy cosmic rays are extremely strong, at the level of $M \gtrsim 10^{10} M_{\rm Pl}$ ($n = 3$, [20]) or $M \gtrsim 10^3$ ($n = 4$ [21]) for protons. In the case $n = 2$ there no energy scale to compare to and constraints are directly on the dimensionless parameter $\Delta c/c$. They are numerically very strong, for example $\Delta c/c_{\gamma e} \lesssim 10^{-13}$ [22, 23].

At these orders in LIV dispersion, neutrinos also are tightly constrained. In this note we review previous constraints on LIV in the neutrino sector, discuss new ones, and detail where the recent OPERA measurement runs into direct confrontation with other constraints.

II. RELEVANT NEUTRINO PROCESSES

The relevant constraints on LIV in neutrino physics for OPERA come from flavor oscillations, time of flight observations, and ultra-high energy neutrino observatories. In this section, we give some background on each process before moving on to constraints.

A. Flavor oscillations

Neutrino flavor oscillations yield constraints on LIV differences within the neutrino sector. Neutrino oscillations depend on the differences in $E-p$ between different neutrino eigenstates. In standard neutrino oscillations, this difference is governed by the squared mass differences between the energy eigenstates. With LIV (and our assumption that the LIV eigenstates are the energy eigenstates) oscillations are governed by the differences in the effective mass squared, denoted $N_i^2 = m_i^2 + \xi_i p^n/M_i^{n-2}$. Therefore, neutrino oscillations do not probe any absolute LIV in the neutrino sector, but rather the differences in any LIV dispersion relations between different neutrino states.

Let us consider a neutrino produced via a particle reaction in a definite flavor eigenstate I with momentum p. The amplitude for this neutrino to be in a particular energy eigenstate i is represented by the matrix U_{Ii}, where $\sum U^\dagger_{Ji} U_{Ii} = \delta_{IJ}$. The amplitude for the neutrino to be observed in another flavor eigenstate J at some distance L from the source, after some time T is then

$$A_{IJ} = \sum_i U^\dagger_{Ji} e^{-i(ET-pL)} U_{Ii} \approx \sum_I U^\dagger_{Ji} e^{-iLN_i^2/(2E)} U_{Ii} \ . \tag{4}$$

The transition probability can then be written as

$$P_{IJ} = \delta_{IJ} - \sum_{i,j>i} 4F_{IJij} \sin^2\left(\frac{\delta N^2_{ij} L}{4E}\right) + 2G_{IJij} \sin^2\left(\frac{\delta N^2_{ij} L}{2E}\right) \ , \tag{5}$$

with $\delta N^2_{ij} = N_i^2 - N_j^2$ and F_{IJij} and G_{IJij} are functions of the mixing matrixes. In the standard formalism used by experimentalists,

$$\delta N^2_{ij} = \Delta m^2_{ij} + p^2 \left(\frac{\Delta c}{c}\right)^{LIV}_{ij} \ . \tag{6}$$

where now

$$\left(\frac{\Delta c}{c}\right)^{LIV}_{ij} = \xi_i \frac{n-1}{2}\left(\frac{p}{M_i}\right)^{n-2} - \xi_j \frac{n-1}{2}\left(\frac{p}{M_j}\right)^{n-2} \tag{7}$$

A nice summary of neutrino oscillation observations, with particular attention to LIV, can be found in [24].

1. *Solar neutrinos*

Neutrinos produced by the Sun at ∼MeV energies yielded the first hint of neutrino oscillations. In the LI framework, their flux can be understood after accounting for oscillations with $\Delta m^2_\odot \simeq 7.58 \times 10^{-5}$ eV^2 [25, 26]. In principle solar neutrinos can also be used to constrain LIV effects. However, in this case the Mikheev-Smirnov-Wolfenstein (MSN) effect [27, 28] must also be taken into account. Since other constraints prove more difficult to reconcile with the OPERA result, we shall ignore solar neutrinos, although in principle the analysis could be done.

2. Atmospheric neutrinos

Muon neutrinos and antineutrinos are produced in interactions of cosmic rays with the Earth atmosphere. Experiments detect preferentially ν_μ and $\bar{\nu}_\mu$. In this case, the survival probability is

$$P_{\nu_\mu,\nu_\mu} \simeq 1 - \sin^2(2\theta_{23}) \sin^2 \left(\frac{\Delta m^2_{atm} L}{4E} + \left(\frac{\Delta c}{c}\right)^{LIV}_{atm} \frac{EL}{4} \right) , \tag{8}$$

where θ_{ij} are the mixing angles of the neutrino mixing matrix. Best fit values (without LIV) are: $\sin^2(\theta_{23}) = 0.42$ and $\Delta m_{atm} \simeq 2.35 \times 10^{-3}$ eV^2 [25, 26].

3. Reactor antineutrinos

Electronic antineutrinos produced by nuclear reactors with $\sim$MeV energy also provide relevant oscillation measurements.

Long baseline The survival probability for long baseline experiments (e.g. KamLAND [29], with a baseline of about 180 km) is

$$P_{\bar{\nu}_e,\bar{\nu}_e} \simeq 1 - \sin^2(2\theta_{12}) \sin^2 \left(\frac{\Delta m^2_{\odot} L}{4E} + \left(\frac{\Delta c}{c}\right)^{LIV}_{\text{long}} \frac{EL}{4} \right) . \tag{9}$$

Best fit values obtained in the standard LI framework are: $\sin^2(\theta_{12}) \simeq 0.3$ and $\Delta m^2_{\odot} = 7.58 \times 10^{-5}$ eV^2 [25, 26].

Short baseline Evidence of electron antineutrino disappearance was sought on much shorter baseline ($L < 1$ km). In this case the survival probability is ruled by a different set of parameters [26]

$$P_{\bar{\nu}_e,\bar{\nu}_e} \simeq 1 - \sin^2(2\theta_{13}) \sin^2 \left(\frac{\Delta m^2_{atm} L}{4E} + \left(\frac{\Delta c}{c}\right)^{LIV}_{\text{short}} \frac{EL}{4} \right) . \tag{10}$$

Experiments reported so far null results so far (i.e. no disappearance), indicating $\theta_{13} \ll 1$ (in facts, hints of $\sin^2(\theta_{13}) \simeq 0.02$ were discussed in [25]). This also means that this observable is not sensitive (yet) to modified kinematics, because the small amplitude of the oscillation is controlled by $\theta_{13} \ll 1$.

4. Accelerator neutrinos

At energy $\gtrsim 1$ GeV a few experiments with short baselines $L \simeq 1$ km have provided evidence for various oscillations, including $\nu_\mu \to \nu_e$, $\nu_\mu \to \nu_\tau$, $\nu_e \to \nu_\tau$ and their conjugates. The T2K Collaboration, with longer baseline, reported recently evidence for oscillation $\nu_\mu \to \nu_e$ [30] at ~ 600 MeV. This process is controlled, as is the short baseline reactor case, by the small angle θ_{13} [26], hence it cannot be used to cast constraints on $(\Delta c/c)^{LIV}$. Also MiniBooNE reported the detection of ν_e. Interestingly, MiniBooNE finds a 3σ excess at 300-500 MeV of $\nu_\mu \to \nu_e$ but only a 1.3σ excess in the conjugate channel [31], hinting perhaps at CPT violation. MiniBooNE also searched for sidereal dependence of the ν_e signal, placing strong constraints on some combination of SME parameters [32].

5. Constraints

The best constraint to date comes from survival of atmospheric muon neutrinos observed by the former IceCube detector AMANDA-II in the energy range 100 GeV to 10 TeV [33], which searched for a generic LIV in the neutrino sector [34] and achieved $(\Delta c/c)_{ij} \leq 2.8 \times 10^{-27}$ at 90% confidence level assuming maximal mixing for some of the combinations i, j. Given that IceCube does not distinguish neutrinos from antineutrinos, the same constraint applies to the corresponding antiparticles. The IceCube detector is expected to improve this constraint to $(\Delta c/c)_{ij} \leq 9 \times 10^{-28}$ in the next few years [35]. The lack of sidereal variations in the atmospheric neutrino flux also yields comparable constraints on some combinations of SME parameters [36].

B. Time of flight

We caution the reader before starting this section: From here on out we attribute a definite velocity to the neutrino flavor eigenstates, although they are not energy eigenstates (or even perhaps LIV eigenstates). However, given the above constraints on the LIV differences between neutrino energy eigenstates, and given that we are considering ultra-relativistic neutrinos, for which the mass term has negligible effect over the CERN to OPERA distance, we can safely refer to the velocity of a flavor eigenstate.

The dispersion relation (3) implies, assuming Hamiltonian dynamics, that the propagation speed of a particle is

$$v(p) \simeq 1 - \frac{m^2}{2p^2} + \xi \left(\frac{n-1}{2}\right)\left(\frac{p}{M}\right)^{n-2} . \tag{11}$$

In turn the time delay (or advance) upon arrival over a path length L, with respect to a light ray traveling at $c \equiv 1$, is

$$\Delta T(p) = \frac{v(p) - c}{c}\frac{L}{c} . \tag{12}$$

We can define

$$\frac{\Delta c}{c}^{TOF}_{(n)} = \frac{v(p) - c}{c} = -\frac{m^2}{2p^2} + \xi \frac{n-1}{2}\left(\frac{p}{M}\right)^{n-2} \tag{13}$$

Given that the standard convention is to include all the factors $(n-1)/2$ into M, we do the same in the following formulae, that can then be derived from Eq. (13) by replacing $M^{n-2} \to M^{n-2}/(n-1)/2$. For $n = 2$, $\Delta c/c$ is a parameter entering directly the modified dispersion relation. Given that $\Delta c/c$ is the quantity "directly" accessible to the experiments at a given energy, we relate the mass scale M to $\Delta c/c$ as

$$M = p \times \left(\frac{\Delta c}{c}^{TOF}_{(n)} + \frac{m^2}{2p^2}\right)^{-1/(n-2)} , \tag{14}$$

where the sign factor ξ has been absorbed in the definition of $\Delta c/c$ to be positive. It is clear from Eq. (12) and Eq. (14) that the constraint placed on M by the measurement of a time delay depends on both the energy and propagation distance.

1. *Constraints*

Unfortunately we have to date only a single astrophysical event for which TOF constraints can be effectively cast on $(\Delta c/c)^{TOF}$. The explosion of SN1987a was a peculiar event which allowed to detect the almost simultaneous (within a few hours) arrival of electronic antineutrinos and photons. Although only a handful of electronic antineutrinos at MeV energies was detected by the experiments KamiokaII, IMB and Baksan, it was enough to establish a constraint $(\Delta c/c)^{TOF} \lesssim 10^{-8}$ [37] or $(\Delta c/c)^{TOF} \lesssim 2\times10^{-9}$ [38] by looking at the difference in arrival time between antineutrinos and optical photons over a baseline distance of 1.5×10^5 ly. Further analyses of the time structure of the neutrino signal, in particular using the fact that the least energetic neutrino in the signal (at 7.5 MeV) was detected within 10 s from the most energetic one (at 30 MeV), strengthened this constraint down to $\sim 10^{-10}$ [39, 40].

The scarcity of the detected neutrino did not allow the reconstruction of the full energy spectrum and of its time evolution. In SN models two main effects are present and lead to the final time-energy structure of the spectrum. First, the SN is globally cooling, hence the average neutrino energy decreases with time; on the other hand, neutrino diffusion in the SN medium depends strongly on the energy of the neutrino, which then determines its escape time. Given these uncertainties, we find constraints purely based on the difference in the arrival time with respect to photons more conservative and robust.

Hence, adopting $\Delta c/c \lesssim 10^{-8}$, the SN constraint implies $M \gtrsim 10^8 \times 10$ MeV $\simeq 10^{-13}$ $M_{\rm Pl}$ for $n = 3$ or $M_\nu \gtrsim$ 0.1 TeV $\simeq 10^{-17}$ $M_{\rm Pl}$ for $n = 4$.

C. Pure LIV effects

LIV introduces, within an EFT description, new processes that a neutrino can undergo. Since neutrinos couple to gravity and photons (via their magnetic moment and charge radius couplings), superluminal neutrinos will emit graviton and photon Čerenkov radiation in vacuum. In addition, high energy superluminal neutrinos will emit neutrino/antineutrino pairs via a neutral current interaction if $n > 2$. The detailed rate computation for these reactions can be found in [20, 41], we merely summarize below.

- Čerenkov radiation in vacuum (photon emission): $\nu \to \nu\gamma$. This possibility has been already investigated for renormalizable operators [43]. Although the rate of this reaction has always been considered too small to produce significant effects even on cosmogenic neutrinos at 10^{20} eV, the effects implied by OPERA are at much lower scales than $M_{\rm Pl}$. Therefore, the rate can be strongly enhanced. The energy loss rate was computed in [20] and found to be

  $$\tau_{\nu\gamma} \simeq \left(\frac{M}{M_{\rm Pl}}\right)^2 \left(\frac{E}{1\ {\rm PeV}}\right)^{-(2n+1)} 10^{26n-86}\ {\rm s}\ . \tag{15}$$

  Gravitational Čerenkov radiation can in principle be considered. However, it is subleading with respect to photon emission and can possibly be an effective energy loss process only for GZK neutrinos with $E \gtrsim 10^{19}$ eV [46]. Furthermore, for $n > 2$ superluminal neutrinos, neutrino splitting dominates both Čerenkov reactions.

- Neutrino splitting ($\nu \to \nu\nu\bar{\nu}$) was studied in the context of ultra-high-energy cosmogenic neutrinos [41]. While the calculation was done for $n = 4$, the methodology is readily adapted to any $n > 2$. In a LI scenario the energy threshold for ν-splitting would be infinite. However, with LIV there is instead a finite energy above which this reaction can happen. The threshold equation for this reaction is analytically solvable when all the neutrinos involved are in the same mass eigenstate, for which the rate is maximal. In this case the energy threshold goes as $E_{th} = (m_\nu^2 M^{n-2})^{1/n}$. If the outgoing neutrino/anti-neutrino pair are in a different eigenstate, then the threshold will change slightly but can still be solved for numerically.

  Using the relation $\xi_{\bar{\nu}} = (-1)^n \xi_\nu$, the typical energy loss time scale for a high energy neutrino with energy well above the threshold energy scales as [2]

  $$\tau_{\nu-{\rm splitting}} \simeq \frac{64\pi^3}{3G_F^2 E^5} \left(\frac{M}{E}\right)^{3(n-2)}\ , \tag{16}$$

  where G_F is the Fermi constant. For example, for the lowest n for which this reaction is allowed, $n = 3$, this corresponds to

  $$\tau \simeq 10^{38} \left(\frac{M}{M_{\rm Pl}}\right)^3 \left(\frac{E_\nu}{10\ {\rm GeV}}\right)^{-8}\ {\rm s}\ . \tag{17}$$

  We caution the reader that there are $O(100)$ phase space factors which vary for each n, so that the lifetime is only approximate. However, since the scaling with M is so strong, these phase factors are largely irrelevant as they change M by only an O(1) factor.

- Neutrino pair production ($\nu \to \nu e^+ e^-$) has been recently proposed in [44]. The calculation was focused on the case $n = 2$ as δ was considered constant when computing the terminal energy for Opera, although much of the calculation and result extends to higher n with only slight modifications. In particular, the rate for neutrino pair production of unmodified electrons will scale the same way with energy as the rate for neutrino splitting far above threshold. The threshold energy will in general depend on n, and can be recovered from a scaling $\delta \to \xi_\nu (E/M)^{n-2}$ only up to O(1) numerical factors.

  The threshold equation reads (for an electron-positron pair of opposite helicity)

  $$\frac{p^n}{M^{n-2}} \left[\xi_\nu^{(n)}(1 - x^{n-1}) - \xi_e(y^{n-1} + (-1)^n t^{n-1})\right] = m_\nu^2 \frac{1-x}{x} + m_e^2 \left(\frac{1}{y} + \frac{1}{t}\right)\ , \tag{18}$$

  where x,y and t are the fraction of initial momentum p carried respectively by the outgoing ν, by e^-, and by e^+, $t = 1 - x - y$ and $0 < x, y, t < 1$.

---

[2] As pointed out by Ward [42], a dimensionless factor of $(E/M)^{n-2}$ was missing in the original rate computed in [41].

The computation of the threshold is straightforward but the general solution is quite cumbersome. Henceforth, we shall provide here only the form of the threshold energy for $n = 3$ and furthermore assume $m_\nu \approx 0$. We also assume no violation in the electron sector $\xi_e = 0$. This is justified because using only the constraints that can be cast on LIV in the electron-positron sector at OPERA energies, any LIV in the electron sector must be much smaller that the one reported by OPERA, at least for $n = 2, 3$.

With these assumptions, the electron/positron pair takes most of the total momentum, so that $x \simeq 0$ at threshold. This implies that the form of the threshold does not depend on n, apart from the scaling of the term p^n/M^{n-2}. As a result the following general scaling holds

$$E^2_{th,(n)} = \frac{4m_e^2}{\delta_{(n)}} \ , \tag{19}$$

with the replacement $\delta_{(n)} = \xi_\nu (E_{th}/M)^{n-2}$.

In addition the rate of this reaction as computed in [44] is general for any n up to numerical factors once one performs the same substitution as before with generic energy $\delta \to \xi_\nu (E/M)^{n-2}$. The generic energy loss time-scale then reads

$$\tau_{\nu-\text{pair}} \simeq G_F^{-2} E^{-5} \left(\frac{M}{E}\right)^{3(n-2)} \ , \tag{20}$$

where we dropped the purely numerical factors. As we see, the rate matches the neutrino splitting rate Eq. (16) up to numerical factors when $n > 2$. The main difference between the two reactions is that pair production is allowed when $n = 2$ while neutrino splitting is expressly forbidden for flavor blind Lorentz breaking.

Pair production alone yields strong constraints on OPERA and so we neglect neutrino splitting as it can only mildly strengthen any constraint. By integrating the energy loss rate from pair production over the distance CERN-Gran Sasso L we obtain

$$E^{-3n+1} - E_0^{-3n+1} = (3n-1)\xi_\nu^3 E_{\text{ref}}^{-3(n-2)} k \frac{G_F^2}{192\pi^3} L \equiv E_T^{-3n+1} \ , \tag{21}$$

where E is the energy on a neutrino starting with energy E_0 after propagation over the distance L and E_{ref} is the energy at which we normalize the parameter ξ_ν. For Opera, $\xi_\nu \sim 5 \times 10^{-5}$ for $E_{\text{ref}} \sim 10$ to 30 GeV. We do this in order to have a direct connection with the observed values. The factor $k = 25/448$ was computed in [44]. The "termination" energy E_T corresponds to the energy that a neutrino would approach after propagation over a distance L if it started with $E_0 \gg E_T$. We remark here that the termination energy E_T is a mildly varying function of n and of the energy scale E_{ref}.

Whereas the oscillation and time of flight supernova constraints are reasonably self-contained, the decay rate constraints are much more tightly tied to the exact characteristics of OPERA and so we devote a larger section to the combined constraints.

III. CAN WE RECONCILE OPERA WITH CURRENT CONSTRAINTS?

A. Energy dependent LIV

It is clear that an obvious problem is to reconcile the OPERA result with the SN constraints, as they are both TOF limits, so they effectively constrain the energy scale at which possible LIV might be found. $\Delta c/c < 10^{-8}$ at ∼10 MeV is incompatible with $\Delta c/c \simeq 10^{-5}$ at ∼10 GeV if LIV is energy independent, as this would be a CPT invariant scenario and LIV differences between various states are so tightly constrained from oscillation data. This alone completely rules out $n = 2$ modified dispersion relation if it has to be valid at all energies, and has additionally been clearly ruled out at OPERA energies alone by [44].

On the other hand, we are also forced to impose $n < 4$. The reason is the following: the OPERA collaboration also measured the neutrino propagation time by dividing the sample of "internal" events, for which it is possible to measure the neutrino energy, in two subsamples with different average energies $\langle E_1 \rangle = 13.9$ GeV, $\langle E_2 \rangle \simeq 42.9$ GeV. No statistically significant differences were found in the time delays of the two samples: $\Delta t_1 = 53.1 \pm 18.8$ (stat.)± 7.4 (sys.),



FIG. 1: Interpretation of time delays with energy dependent LIV. Blue: $n = 4$ LIV, with $M = 3.5$ TeV. Red: $n = 3$ LIV. Solid: lines drawn to reproduce the time advance obtained with the full sample of average energy 17 GeV (and with $M = 700$ TeV for the $n = 3$ case. Dashed: attempt of a fit with $n = 3$ that might reproduce also the measurement at the 2 different energies, with $M = 10^3$ TeV. In the $n = 4$ case the fit is basically impossible. Errors on the x-axis are inferred from the characteristics of the sample, not given by the experimental Collaboration.

$\Delta t_2 = 67.1 \pm 18.2\,(\text{stat.}) \pm 7.4\,(\text{sys.})$. If we now compute the time delay on a baseline of ~ 730 km assuming $n = 4$ we find time delays incompatible with the observations, as shown in Fig. 1.

Therefore, we are left only with $n = 3$, which, according to Fig. 1, can possibly explain all the time of flight constraints assuming $M \sim 10^{-13}\ M_{\rm Pl}$. The agreement with OPERA data is however achieved only at the 2σ level. A better agreement might be achieved if we could convolve the energy dependent TOF prediction with the actual OPERA neutrino spectrum. Notice that, as we mentioned in the Introduction, the case $n = 3$ includes CPT odd LIV, which implies $\xi_\nu = -\xi_{\bar\nu}$. Therefore superluminal neutrinos imply subluminal antineutrinos. If moreover we assume flavor independence, we do not alter flavor oscillations. In the $n = 3$ case, the absence of significant energy dependence is interpreted as due to poor statistics. It is an easy exercise to verify that in the $n = 3$ case the SN constraint $\Delta c/c \lesssim 10^{-8}$ is respected.

However, superluminal neutrinos can possibly emit e^+e^- pairs, or they can split according to $\nu \to \nu\nu\bar\nu$ (since antineutrinos are necessarily subluminal), thereby losing energy. According to Eq. (21), in the case $n = 3$ the argument by [44] applies albeit with $E_{th} \simeq 1.5$ GeV and terminal energy $E_T \simeq 15$ GeV, which again rules out at OPERA energies the most mildly energy dependent dispersion relation $n = 3$.

Furthermore, according to Eq. (16), atmospheric muon neutrinos at 10 TeV would have a typical mean free path of about 3×10^{-15} cm (the mean free path implied by Čerenkov is 3×10^{-14} cm, see Eq. 15) for $M \sim 10^{-13}\ M_{\rm Pl}$, too short to be compatible with the observation of upward-going atmospheric neutrinos up to 400 TeV by IceCube. [3]

Therefore, LIV in EFT with any energy dependence does not seem a viable scenario.

B. Flavor dependent LIV

Given that the flavor independent LIV scenario has problems, one might be tempted to increase the degrees of freedom of the problem and reconsider a form of LIV that depends on flavor. In this way, the SN constraint does not need be necessarily respected by OPERA neutrinos, because it is cast on $\bar\nu_e$, while OPERA observes ν_μ. However, the constraints on $n = 2$ [44] and on $n = 3$ just seen in the previous section still apply also in the flavor dependent case. Hence flavor dependence cannot be considered a solution.

Nonetheless, we want to briefly show here that even forgetting the above mentioned constraints, it is still possible to rule out both $n = 2$ and $n = 3$ possibilities simply on the basis of neutrino flavor oscillations.

First of all, given the atmospheric neutrino constraints reported in section (II A 5), we have to demand that the

---

[3] While submitting this paper we found that a similar remark was also made in [45].



FIG. 2: Survival probability $P_{\bar{\nu}_e,\bar{\nu}_e}$ for different values of $(\Delta c/c)^{LIV}$. The red solid curve is the standard LI case with solar neutrino parameters. Green dotted and black dashed curves are for $(\Delta c/c)^{LIV} = 10^{-16}$ and 5×10^{-16} respectively.

LIV parameters be closer than 10^{-27} for $n = 2$, or 10^{-31} for $n = 3$ (see Eq. 7) if $M \simeq 10^{-13}\ M_{\text{Pl}}$ for $\nu_\mu - \nu_\tau$ and $\bar{\nu}_\mu - \bar{\nu}_\tau$ oscillations.

We now consider reactor antineutrinos. For our purpose we can use Eq. (9) to understand what happens in the LIV case. In Fig. 2 we show how even a tiny modification $\Delta c/c \simeq 10^{-16}$ would lead to large unacceptable departures from the LI case. Indeed, KamLAND observed [29] the first 2 humps of the survival probability at values $L/E \sim 30$ to 50 km/MeV, in clear disagreement with the 4 humps present for $\Delta c/c \sim 10^{-16}$. Given the TOF constraints, in the computation of LIV effects on oscillations we have to require $(\Delta c/c)^{LIV}_{\text{long}} \simeq 10^{-5}$ for $n = 2$ and $(\Delta c/c)^{LIV}_{\text{long}} \simeq 10^{-8}$ for the CPT odd $n = 3$ scenario at MeV energies. Hence, these values are far too large to be compatible with the observed oscillations. Therefore a flavor dependent LIV case is strongly disfavored.

IV. CONCLUSIONS

In this note we discussed the OPERA results. Our conclusions, in agreement and complementing other similar analyses [44, 47, 48], are that these results are not compatible with other observations and derivable constraints within an EFT framework (even models predicting modified dispersion relations on finite energy ranges, including OPERA energies [48]).[4]

Of course, an experiment is not proved wrong by the theoreticians' inability to come up with viable explanations, and the last word on the reliability of the OPERA measurement lays in the hands of forthcoming experiments aimed at reproducing the results. However, we think that taken at face value, this analysis (and the above mentioned similar ones) should be taken as a strong case for being cautious in developing freely new exotic models aimed at fitting OPERA data, or at least as guidance about what these models should actually provide.

It is difficult to envisage which models can escape the wealth of constraints at these energies, in particular the one provided by the absence of pair production by OPERA neutrinos. One example could be models like Deformed Special Relativity (DSR) which do not lead to new threshold reactions [53]. [5]

Alternatively, one can envisage the possibility that in vacuo neutrino propagation is not affected itself by LIV, while some LIV interaction can be the source of OPERA anomaly. In the past there have been models, most noticeably the so called space-time foam ones, like [55], predicting a modified propagation only in some SM sector (photons and neutrinos in [55]) and only on average, not instantaneously, due to many interactions with some hidden sector of the theory (in the case of [55], string theory D-branes). Averaging over many of these interaction events leads to features not predictable via standard EFT reasoning. However the model described in [55] predicts subluminal propagation of photons and neutrinos [6], and hence it is not amenable to explain Opera results, although it is possible that different implementations of the idea that lead to superluminal neutrinos may yield a workable proposal.

---

[4] See also [49, 50] for alternative albeit less stringent arguments. For tentative explanations see e.g. [51, 52].

[5] Unfortunately, the various DSR candidates face in general major problems regarding their physical interpretation and a working model is not yet available (see however [54] for recent attempts in new directions).

Models of this kind could possibly evade most if not all the constraints based on particle interactions (see however [56] for a constraint on a possible incarnation of the framework discussed in [55]) and still lead to modified propagation on sufficiently long distances. Furthermore, the "hidden" fields can be homogeneously distributed in the Universe, or highly clumped, e.g. around the Earth's position. This could perhaps play an important role, like for example in the model of [57]. If OPERA is confirmed, there is a long road ahead for model builders trying to match all the existing constraints. However, precisely because of this wealth of data we have, there is a lot of experimental guidance on how one should proceed.

Acknowledgments

We are grateful to Ted Jacobson for useful remarks and suggestions. LM gratefully thanks Alessandro Mirizzi for enlightening discussions on neutrino physics. SL wishes to thank Matt Visser and Marco Serone for useful remarks, references and discussions. LM acknowledges support from the State of Hamburg, through the Collaborative Research program "Connecting Particles with the Cosmos" within the framework of the LandesExzellenzInitiative (LEXI). DM thanks the University of New Hampshire for research support and Matt Reinhold for useful discussions.

[1] T. Adam *et al.* [OPERA Collaboration], [arXiv:1109.4897 [hep-ex]].

[2] M. Antonello *et al.* [ICARUS Collaboration], arXiv:1203.3433 [hep-ex].

[3] P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. **D76** (2007) 072005. [arXiv:0706.0437 [hep-ex]].

[4] A. Kogut, C. Lineweaver, G. F. Smoot, C. L. Bennett, A. Banday, N. W. Boggess, E. S. Cheng, G. De Amici *et al.*, Astrophys. J. **419** (1993) 1. [astro-ph/9312056].

[5] R. Gambini and J. Pullin, Phys. Rev. D **59** (1999) 124021 [gr-qc/9809038].

[6] C. P. Burgess, J. M. Cline, E. Filotas, J. Matias and G. D. Moore, JHEP **0203**, 043 (2002) [hep-ph/0201082].

[7] S. M. Carroll, J. A. Harvey, V. A. Kostelecky, C. D. Lane and T. Okamoto, Phys. Rev. Lett. **87**, 141601 (2001) [hep-th/0105082].

[8] C. Barcelo, S. Liberati and M. Visser, Living Rev. Rel. **8**, 12 (2005) [gr-qc/0505065].

[9] A. Kostelecky and M. Mewes, arXiv:1112.6395 [hep-ph].

[10] V. A. Kostelecky and M. Mewes, Phys. Rev. D **70**, 031902 (2004) [hep-ph/0308300].

[11] O. W. Greenberg, Phys. Rev. Lett. **89** (2002) 231602. [hep-ph/0201258].

[12] J. Collins, A. Perez, D. Sudarsky, L. Urrutia and H. Vucetich, Phys. Rev. Lett. **93** (2004) 191301 [gr-qc/0403053].

[13] R. Gambini, S. Rastgoo and J. Pullin, Class. Quant. Grav. **28** (2011) 155005 [arXiv:1106.1417 [gr-qc]].

[14] J. Polchinski, arXiv:1106.6346 [gr-qc].

[15] T. Jacobson, S. Liberati, D. Mattingly, Annals Phys. **321** (2006) 150-196. [astro-ph/0505267].

[16] P. Laurent, D. Gotz, P. Binetruy, S. Covino, A. Fernandez-Soto, [arXiv:1106.1068 [astro-ph.HE]].

[17] F. W. Stecker, Astropart. Phys. **35**, 95-97 (2011). [arXiv:1102.2784 [astro-ph.HE]].

[18] S. Liberati, L. Maccione, Ann. Rev. Nucl. Part. Sci. **59** (2009) 245-267. [arXiv:0906.0681 [astro-ph.HE]].

[19] L. Maccione, S. Liberati, A. Celotti, J. G. Kirk, JCAP **0710** (2007) 013. [arXiv:0707.2673 [astro-ph]].

[20] T. Jacobson, S. Liberati, D. Mattingly, Phys. Rev. **D67** (2003) 124011. [hep-ph/0209264].

[21] L. Maccione, A. M. Taylor, D. M. Mattingly, S. Liberati, JCAP **0904** (2009) 022. [arXiv:0902.1756 [astro-ph.HE]].

[22] D. Mattingly, Living Rev. Rel. **8** (2005) 5. [gr-qc/0502097].

[23] V. A. Kostelecky, N. Russell, Rev. Mod. Phys. **83** (2011) 11. [arXiv:0801.0287 [hep-ph]].

[24] J. S. Diaz, A. Kostelecky, [arXiv:1108.1799 [hep-ph]].

[25] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo, A. M. Rotunno, [arXiv:1106.6028 [hep-ph]].

[26] K. Nakamura et al. (Particle Data Group), J. Phys. G **37** (2010) 075021 and 2011 partial update for the 2012 edition.

[27] L. Wolfenstein, Phys. Rev. **D17** (1978) 2369-2374.

[28] S. P. Mikheev, A. Y. .Smirnov, Sov. J. Nucl. Phys. **42** (1985) 913-917.

[29] S. Abe *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **100** (2008) 221803. [arXiv:0801.4589 [hep-ex]].

[30] K. Abe *et al.* [T2K Collaboration], Phys. Rev. Lett. **107** (2011) 041801. [arXiv:1106.2822 [hep-ex]].

[31] A. A. Aguilar-Arevalo *et al.* [MiniBooNE Collaboration], Phys. Rev. Lett. **102** (2009) 101802. [arXiv:0812.2243 [hep-ex]].

[32] The MiniBooNE Collaboration, [arXiv:1109.3480 [hep-ex]].

[33] J. L. Kelley [IceCube Collaboration], Nucl. Phys. **A827** (2009) 507C-509C.

---

[6] Note however that in this model LIV is present for Majorana neutrinos, while we considered here only Dirac-type neutrinos.

[34] M. C. Gonzalez-Garcia, M. Maltoni, Phys. Rev. **D70** (2004) 033010. [hep-ph/0404085].
[35] W. Huelsnitz *et al.* [IceCube Collaboration], Proceedings of the 31st ICRC.
[36] R. Abbasi *et al.* [IceCube Collaboration], Phys. Rev. **D82** (2010) 112003. [arXiv:1010.4096 [astro-ph.HE]].
[37] L. Stodolsky, Phys. Lett. **B201** (1988) 353.
[38] M. J. Longo, Phys. Rev. Lett. **60** (1988) 173.
[39] J. R. Ellis, N. Harries, A. Meregaglia, A. Rubbia, A. Sakharov, Phys. Rev. **D78** (2008) 033013. [arXiv:0805.0253 [hep-ph]].
[40] A. Sakharov, J. Ellis, N. Harries, A. Meregaglia, A. Rubbia, J. Phys. Conf. Ser. **171** (2009) 012039. [arXiv:0903.5048 [hep-ph]].
[41] D. M. Mattingly, L. Maccione, M. Galaverni, S. Liberati, G. Sigl, JCAP **1002** (2010) 007. [arXiv:0911.0521 [hep-ph]].
[42] B. F. L. Ward, arXiv:1201.1322 [hep-ph].
[43] S. R. Coleman, S. L. Glashow, Phys. Rev. **D59** (1999) 116008. [hep-ph/9812418].
[44] A. G. Cohen, S. L. Glashow, [arXiv:1109.6562 [hep-ph]].
[45] J. M. Carmona, J. L. Cortes, [arXiv:1110.0430 [hep-ph]].
[46] J. Alexandre, J. Ellis, N. E. Mavromatos, [arXiv:1109.6296 [hep-ph]].
[47] G. Amelino-Camelia, G. Gubitosi, N. Loret, F. Mercati, G. Rosati, P. Lipari, [arXiv:1109.5172 [hep-ph]].
[48] G. F. Giudice, S. Sibiryakov, A. Strumia, [arXiv:1109.5682 [hep-ph]].
[49] X. -J. Bi, P. -F. Yin, Z. -H. Yu, Q. Yuan, [arXiv:1109.6667 [hep-ph]].
[50] B. Altschul, [arXiv:1110.2123 [hep-ph]].
[51] C. Pfeifer, M. N. R. Wohlfarth, [arXiv:1109.6005 [gr-qc]].
[52] M. Matone, [arXiv:1109.6631 [hep-ph]].
[53] G. Amelino-Camelia, Symmetry **2** (2010) 230-271. [arXiv:1003.3942 [gr-qc]].
[54] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman, L. Smolin, [arXiv:1101.0931 [hep-th]].
[55] J. Ellis, N. E. Mavromatos, D. V. Nanopoulos, Int. J. Mod. Phys. **A26** (2011) 2243-2262. [arXiv:0912.3428 [astro-ph.CO]].
[56] L. Maccione, S. Liberati, G. Sigl, Phys. Rev. Lett. **105**, 021101 (2010). [arXiv:1003.5468 [astro-ph.HE]].
[57] G. Dvali, A. Vikman, [arXiv:1109.5685 [hep-ph]].